

A LIMIT RELATION BETWEEN BLACK HOLE MASS AND H β WIDTH: TESTING SUPER-EDDINGTON ACCRETION IN ACTIVE GALACTIC NUCLEI

JIAN-MIN WANG^{1,2}

¹Institut für Astronomie und Astrophysik, Abt. Astronomie, Universität Tübingen, Sand 1, D-72076 Tübingen, Germany

²Laboratory for High Energy Astrophysics, Institute of High Energy Physics, CAS, Beijing 100039, P.R. China
Astronomical Journal, vol. 125 (June Issue, 2003), in press

ABSTRACT

In this paper, we show that there is a limit relation between the black hole mass (M_{BH}) and the width at the half maximum (v_{FWHM}) of H β for active galactic nuclei (AGNs) with super-Eddington accretion rates. When a black hole has a super-Eddington accretion rate, the empirical relation of reverberation mapping has two possible ways. First, it reduces to a relation between the black hole mass and the size of the broad line region due to the photon trapping effects inside the accretion disk. For the Kaspi et al.'s empirical reverberation relation, we get the limit relation as $M_{\text{BH}} = (2.9 - 12.6) \times 10^6 M_{\odot} (v_{\text{FWHM}}/10^3 \text{ km s}^{-1})^{6.67}$, called as the Eddington limit. Second, the Eddington limit luminosity will be relaxed if the trapped photons can escape from the magnetized super-Eddington accretion disk via the photon bubble instability, and the size of the broad line region will be enlarged according to the empirical reverberation relation, leading to a relatively narrow width of H β . We call this the Begelman limit.

Super-Eddington accretions in a sample composed of 164 AGNs have been searched by this limit relation. We find that most of them are well confined by the Eddington limit relation, namely most of the objects in the sample have sub-Eddington accretion rates, but there are a handful of objects locate between the Eddington and Begelman limit lines, they may be candidates of super-Eddington accretors in a hybrid structure of photon trapping and photon bubble instability. The maximum width of H β is in the range of $(3.0 - 3.8) \times 10^3 \text{ km s}^{-1}$ for the maximum mass black holes with super-Eddington accretion rates among AGNs. We suggest that the FWHM(H β)– M_{BH} relation is more reliable and convenient to test whether a source is super-Eddington and useful to probe the structure of the super-Eddington accretion process.

Subject headings: accretion: super-Eddington - active galaxies - black hole

1. INTRODUCTION

Accretion onto a supermassive black hole is regarded as the energy sources of AGNs (Rees 1984). Energy output from accretion disks strongly depends on their structures and radiation efficiency, which is mainly controlled by the Eddington ratio, defined by the dimensionless accretion rate $\dot{m} = \dot{M}/\dot{M}_{\text{Edd}}$, where $\dot{M}_{\text{Edd}} = 1.39 \times 10^{18} \eta_{-1}^{-1} (M_{\text{BH}}/M_{\odot}) \text{ g s}^{-1}$, $\dot{M}$ is the accretion rate, M_{BH} the black hole mass and the accretion efficiency $\eta = 0.1\eta_{-1}$ (Chen et al. 1997, Frank, King & Raine 2002). It has been realized that many observable features of AGNs could be driven by $\dot{m}$ (Boroson & Green 1992, Boroson 2002), however, it is not easy to estimate $\dot{m}$ from observations.

We need a more reliable and convenient criteria based on theory to test whether AGNs have super-Eddington accretion rates. Much attention has been given to estimate the black hole masses in AGNs and inactive galaxies (Kormendy & Gebhardt 2001) so that the black hole mass becomes an easily "measured" quantity. There are currently four fundamental ways to reliably estimate the mass of the black holes by: 1) reverberation mapping methods for AGNs (Wandel, Peterson & Malkan 1999, Kaspi et al. 2000), 2) the empirical relation of the reverberation mapping (Netzer 2003, Vestergaard 2002), 3) relation between $M_{\text{R}} - M_{\text{BH}}$ (Kormendy & Richstone 1995, Ho 1999, McLure & Dunlop 2001), where M_{R} is the absolute R-band magnitudes of the host galaxies, 4) the relation $\sigma - M_{\text{BH}}$ (Ferrarese & Merrit 2000, Gebhardt et al. 2000, Tremaine et al. 2002), including the method from the fundamental plan, where σ is the dispersion velocity. Employing the empirical relation of reverberation mapping, one can conveniently obtain the masses of the black holes in a large sample spanning a wide range of redshift (Netzer 2003). With these estimations, the Ed-

dington ratio can be estimated via $L/L_{\text{Edd}} = \xi L_{5100}/L_{\text{Edd}}$, where $\xi = 5 - 9$ (Kaspi et al. 2000, Netzer 2003). It has been found that some AGNs may have super-Eddington accretion rates due to $L/L_{\text{Edd}} > 1$. They consist of some narrow line Seyfert 1 galaxies and quasars (Collin & Huré 2001, Collin et al. 2002, King & Puchnarewicz 2002, Wang & Netzer 2003). One of the most prominent features is the strong soft X-ray humps (King 2002, King & Puchnarewicz 2002, Wang & Netzer 2003).

The structure of the super-Eddington accretion disk remains open. Begelman (1978) and Begelman & Meier (1982) suggested that a photon trapping process may happen in a super-Eddington accretion. The classical model of the slim accretion disk ($\dot{m} < 50 - 100$) based on the vertical averaged equations advocated by Abramowicz et al (1988) removes the thermal instability of the radiation pressure-dominated region since the advection as an efficient cooling dominates over the diffusive cooling from the disk surface (Chen & Taam 1993). The detail treatment of the photon trapping inside the slim disk has been done by Ohsuga et al. (2002), who showed the maximum luminosity is about Eddington luminosity. This is roughly in agreement with Wang & Zhou (1999). On the other hand, Gammie (1998) showed that a *magnetized* accretion disk may have photon bubble instability in the radiation pressure-dominated region of the standard accretion disk model (Shakura & Sunyaev 1973). Such an instability in the super-Eddington accretion disks should be enhanced since the photons are trapped inside the disks and the radiation pressure is much higher than that in Shakura & Sunyaev's disk. Begelman (2002) suggested that the inhomogeneous distribution of the accreted matter will be formed due to the photon bubble instability and the maximum luminosity will exceed Eddington luminosity by a factor

of one hundred for galactic mass of the black hole. Both the spectra and the behavior of variabilities from slim disk (Wang et al. 1999; Wang & Netzer 2003; Mineshige et al. 2000) are not sufficiently understood as well as the roles of the photon bubble instability (Begelman 2002). It may be difficult to directly test the structure of the super-Eddington accretion disk in an individual objects, but statistic test may provide invaluable information.

We show in this paper that $L/L_{\text{Edd}} = \xi L_{5100}/L_{\text{Edd}}$ can not properly represent the accretion disk rates in AGNs, particularly for those with super-Eddington accretion rates. We suggest a limit relation between the black hole mass and $H\beta$ width based on the theoretical models to test whether the accretion disks in AGNs have super-Eddington accretion rates. These limit relations are useful to probe the structures of the disks with super-Eddington rates.

2. THE EDDINGTON AND BEGELMAN CONSTRAINTS

2.1. Size of the broad line region

Measuring the response of the broad emission line to the variations of the continuum can provide the distance of the clouds from the central black hole in AGNs. Kaspi et al. (2000) provided the results of the reverberation mapping in a sample consisting of 34 AGNs and found a strong correlation between the size of the broad line region (R_{BLR}) and the luminosity at 5100Å, namely, the empirical reverberation relation

$$R_{\text{BLR}} = \mathcal{R}_0 \left(\frac{\lambda L_\lambda}{10^{44} \text{erg s}^{-1}} \right)^\beta \text{ lt-days}, \quad (1)$$

where $\beta = 0.7$, $\mathcal{R}_0 = 32.9$ and λL_λ is the continuum luminosity at wavelength $\lambda = 5100\text{\AA}$. This relation has been calibrated by Vestergaard (2002) using the bivariate correlated errors and intrinsic scatter algorithm. The specific values of $\mathcal{R}_0$ and β are in debate (see Vestergaard 2002, Netzer 2003). A constant ionization parameter U , defined by $U = L_{\text{ion}}/4\pi R^2 N_e$, leads to a relation $R_{\text{BLR}} \propto L_{\text{ion}}^{0.5}$ (Wandel, Peterson & Malkan 1999), where N_e is the number density of electrons in ionized medium and L_{ion} is the ionizing luminosity. This is supported by the argument that the outer boundary of the BLR is determined by dust sublimation radius (Netzer & Laor 1993). We keep the uncertainty of the index β in mind, however it does not influence the existence of the limit relation between M_{BH} and $H\beta$ width derived in the present paper.

For a given black hole, the continuum luminosity from an accretion disk increases with the accretion rate (but below the Eddington rate) and hence an increase of the BLR size according to the empirical reverberation relation (equation 1). This implies that a high accretion rate leads to a relatively narrow width of $H\beta$ for a sub-Eddington accretion disk in AGN if the assumption that the broad line emitting-clouds are virialized in the black hole potential works (see equation 2). When the accretion rate exceeds the Eddington rate, the radiated luminosity from the disk very weakly depends on the accretion rate ($L_{\text{disk}} \propto \log \dot{M}$), but linearly proportional to the mass of the black hole in the context of the slim disk model (Abramowicz et al. 1988, Wang & Zhou 1999; Mineshige et al. 2000; Ohsuga et al. 2002). Thus the radiated luminosity is *saturated*, leading to a *constant* BLR size independent of the accretion rate for a fixed black hole. We call this the Eddington size and the Eddington width of the emission line accordingly. For AGNs with super-Eddington accretion rates, the Eddington width of $H\beta$ directly reflects the potential of the black holes. In the context of

the photon trapping disk a relation of M_{BH} and $\text{FWHM}(H\beta)$ is then expected.

However, it has been argued that magnetized disks with dominance of radiation pressure are suffering from photon bubble instability (Gammie 1998). The disks become so inhomogeneous on scales much smaller than the radiation pressure scale height that the radiation will be squeezed by gravity out of overdense region enhanced by the magnetic tension while the underdense regions become tenuous due to the acceleration of radiation force. Some of the trapped photons may be liberated from the disk via this instability. Thus the Eddington limit may be relaxed by this way. Begelman (2001, 2002) investigated the photon bubble instability in a super-Eddington accretion disk and found the maximum radiated luminosity $L_{\text{max}} \approx 300 M_{\text{BH},6}^{1/5} L_{\text{Edd}}$, where $M_{\text{BH},6} = M_{\text{BH}}/10^6 M_\odot$, keeping a geometrically thin shape, when the accretion rate reaches its maximum. We call this the Begelman limit. It is not clear what the radiated luminosity is before $\dot{M}$ reaches its maximum as well as the emergent spectrum from such disks. If the photon bubble instability works as a mechanism liberating the trapped photons in AGN super-Eddington accretion disks, the BLR size will be larger than the Eddington size, resulting in a narrower width of $H\beta$ than the Eddington width. Thus the different structure of the super-Eddington accretion disk should lead to different limit relations between $\text{FWHM}(H\beta)$ and M_{BH} . This lends us a possibility to statistically test the structure of the super-Eddington accretion disk based on the available data of $\text{FWHM}(H\beta)$ and M_{BH} .

2.2. A limit relation between $\text{FWHM}(H\beta)$ and M_{BH}

The broad line $H\beta$ emitting clouds are virialized in the gravitational potential of the supermassive black hole, the mass of the black hole is expressed by the $\text{FWHM}(H\beta)$

$$M_{\text{BH}} = 1.46 \times 10^5 f^2 \left(\frac{R_{\text{BLR}}}{\text{lt-days}} \right) \left(\frac{v_{\text{FWHM}}}{10^3 \text{km s}^{-1}} \right)^2 M_\odot, \quad (2)$$

where we take $v = \sqrt{3} f v_{\text{FWHM}}/2$ for the corrections of geometry and kinetics (Peterson & Wandel 1999, Fromerth & Melia 2000, McLure & Dunlop 2001, Krolik 2001). The factor f remains a matter for debate of BLR geometry, but $f = 1$ is expected. We need to calculate the luminosity at 5100Å from the emergent spectrum from the slim disk. This has been done in detail by Wang et al. (1999). We find that $\varepsilon = L_{\text{disk}}/L_{5100} = 4.0 - 7.5$ for a supermassive black hole. Combining the equations (1) and (2), we have a relation between $\text{FWHM}(H\beta)$ and M_{BH} ,

$$M_{\text{BH},6} = \left(\frac{\mathcal{E}_R}{C} \right)^{\beta/(1-\beta)} v_{\text{FWHM},3}^{2/(1-\beta)}, \quad (3)$$

where $v_{\text{FWHM},3} = v_{\text{FWHM}}/10^3 \text{km s}^{-1}$ and the constant C is

$$C = \frac{\varepsilon}{1.26} \left(\frac{10}{1.46 \mathcal{R}_0 f^2} \right)^{1/\beta}, \quad \text{and} \quad \mathcal{E}_R = \frac{L_{\text{disk}}}{L_{\text{Edd}}}. \quad (4)$$

This relation clearly relies on two model-dependent parameters ε and $\mathcal{E}_R$, which are determined by the spectrum and the structure of the super-Eddington accretion disk. Such a dependence can thus be used to probe the structures of the super-Eddington accretion disks.

The photon trapping process efficiently lowers the radiated luminosity from the disk, resulting in a very weak dependence

of the radiated luminosity on $\log \dot{M}$. For a photon trapping accretion flow, $\mathcal{E}_R \approx 1$ (Wang & Zhou 1999, Ohsuga et al. 2002), we have the Eddington limit from equation (3) as

$$M_{\text{BH},6} = C_1 v_{\text{FWHM},3}^{2/(1-\beta)}, \quad (5)$$

where $C_1 = C\beta/(\beta-1)$. For the empirical relation of reverberation mapping given by Kaspi et al. (2000), $\beta = 0.7$ and $\mathcal{R}_0 = 32.9$, we have a limit relation of $M_{\text{BH},6} = C_1 v_{\text{FWHM},3}^{6.67}$, where $C_1 = 2.9 - 12.62$ for $\varepsilon = 4.0 - 7.5$ and $f = 1$. The Eddington limit lines are shown in Figure 1. This is a very strong dependence on $\text{FWHM}(\text{H}\beta)$. Those objects below this line in the $\text{FWHM}(\text{H}\beta) - M_{\text{BH}}$ plot have sub-Eddington accretion rates.

On the other hand, the photon bubble instability may operate in a super-Eddington accretion disk and the maximum luminosity from the disk reads $L_{\text{max}} \approx \mathcal{E}_{R,\text{max}} M_{\text{BH},6}^{1/5} L_{\text{Edd}}$ where $\mathcal{E}_{R,\text{max}} \approx 300$ for the maximum accretion rate (Begelman 2002). Such a high luminosity will strongly affect the BLR size, equation (3) for the Begelman limit is changed to

$$M_{\text{BH},6} = C_2 v_{\text{FWHM},3}^{10/(5-6\beta)} \quad (6)$$

where $C_2 = (\mathcal{E}_{R,\text{max}}/C)^{5\beta/(5-6\beta)}$. The spectrum from a disk with the photon bubble instability is highly unknown, we take $\varepsilon = 4$ to get C_2 for an illustration, we have $M_{\text{BH},6} = 8.0 \times 10^{12} v_{\text{FWHM},3}^{12.5}$ if the Kaspi et al.'s relation is used. This relation is highly different from that determined by the photon trapping process. For a disk with the photon bubble instability, the $\text{FWHM}(\text{H}\beta) - M_{\text{BH}}$ relation is beyond the scope of the frame of Figure 1, we draw $\mathcal{E}_R = 2$ line for the illustration of the Begelman limit in Figure 1. A potentially realistic situation may be a hybrid structure undergoing the photon trapping and photon bubble instability in the region between the Eddington and the Begelman lines, where some objects appear if they have a super-Eddington accretion rate.

The Eddington ratio is usually estimated by $L/L_{\text{Edd}} = \xi L_{5100}/L_{\text{Edd}}$, where $\xi = 5 - 9$ (Kaspi et al. 2000, Telfer et al. 2002, Netzer 2003). If $L/L_{\text{Edd}} > 1$, this object is generally thought to be a super-Eddington accretor. However, we argue that $L/L_{\text{Edd}} = \xi L_{5100}/L_{\text{Edd}} > 1$ does not simply imply a super-Eddington accretion in practice. If an object with $\xi L_{5100}/L_{\text{Edd}} > 1$ really has a super-Eddington accretion rate, it should locate on the Eddington limit line (photon trapping), or between the Begelman and Eddington lines (a hybrid structure of photon trapping and photon bubble instability). The $\text{FWHM}(\text{H}\beta) - M_{\text{BH}}$ relation can be used to reliably test the super-Eddington accretion in AGNs.

It should be noted that the $\text{FWHM}(\text{H}\beta) - M_{\text{BH}}$ relation is very sensitive to β due to the indices $2/(1-\beta)$ and $10/(5-6\beta)$. Any small calibrations on the empirical relation of reverberation mapping will be important to the limit relation. $\beta = 0.58$ and $\mathcal{R}_0 = 33.1$ were suggested by Netzer (2003) for the cosmology assumed in the present paper $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.3$ and $\Omega_\Lambda = 0.7$ based on the BCES (bivariate correlated errors and intrinsic scatter) estimator. We plot the Eddington limit lines for both of the Kaspi et al.'s and Netzer's values of $(\beta, \mathcal{R}_0)$, and the Begelman limit line for Kaspi et al.'s value in Figure 1.

3. SUPER-EDDINGTON ACCRETIONS IN AGNS

It is interesting to search super-Eddington accretions and to test their structures in AGNs via the limit relation of $\text{FWHM}(\text{H}\beta) - M_{\text{BH}}$. We assemble objects from the literatures,




FIG. 1. — The plot of $v_{\text{FWHM}}(\text{H}\beta) - M_{\text{BH}}$. The dotted and solid lines labelled by the Eddington limit are from equation (5) for $(\beta = 0.58, \mathcal{R}_0 = 33.1)$ and $(\beta = 0.7, \mathcal{R}_0 = 32.9)$. The different color symbols represent the different samples. The open circles represent the sources with $L/L_{\text{Edd}} < 1$ and filled circles those with $L/L_{\text{Edd}} > 1$. Eddington limit lines (equation 5) are labeled by $\varepsilon = 4$ and $\varepsilon = 7.5$. Begelman limit line is drawn for $\mathcal{E}_R = 2$ for illustration. The objects labelled by numbers are listed in Table 1 for Kaspi et al.'s relation.

in which their black hole masses can be estimated by 1) the reverberation mapping, 2) the $M_R - M_{\text{BH}}$ relation and 3) the $\sigma - M_{\text{BH}}$ relation. The three methods are independent of the method of the empirical reverberation relation as used by Netzer (2003) in a large sample of AGNs. This allows us to test the super-Eddington accretion disks in AGNs. We still estimate the Eddington ratio via $L/L_{\text{Edd}} = \xi L_{5100}/L_{\text{Edd}}$, where $\xi = 5$ (Telfer et al. 2002, Netzer 2003) in order to compare with the $\text{FWHM}(\text{H}\beta) - M_{\text{BH}}$ limit relation.

We omitted five objects (IC 4329A, NGC 3227, NGC 7469, PG 1700 and PG 1704) in Kaspi et al.'s sample since their lower uncertainties are too large. There are 30 objects measured by reverberation mapping, 29 from Kaspi et al.'s sample and NGC 3783 from Onken & Peterson (2002). McLure & Dunlop (2001) present 45 objects with host galaxy absolute magnitude M_R , in which 15 Seyfert galaxies of objects have been measured by reverberation mapping (Kaspi et al. 2000). We thus use the results of the reverberation mapping but we take the absolute magnitudes of IC 4329A, NGC 3227 and NGC 7469 since their lower limits of the black hole mass are too uncertain by the reverberation mapping. We take 17 of 37 radio-loud quasars with M_R assembled by Wang, Ho & Staubert (2002) [the rest radio-loud quasars in Wang et al. (2002) overlap with that given by McLure & Dunlop (2001)]. We then use the relation $\log(M_{\text{BH}}/M_\odot) = -0.5M_R - 2.96$ to get the masses of the black holes (McLure & Dunlop 2002). Shields et al. (2002) extend the relationship $\sigma - \text{FWHM}([\text{O III}])$ in Nelson (2000). Using $\text{FWHM}([\text{O III}])$ as the stellar dispersion velocity σ , the relation $M_{\text{BH}} = 10^{8.13} (\sigma/200 \text{ km s}^{-1})^{4.02} M_\odot$ (Tremaine et al. 2002) enables us to reliably estimate the masses of the black holes in 84 quasars (35 RLQs + 49 RQQs) in Shields' sample. The error bars in this sample are given by the largest intrinsic scatter ($< 30\%$) (Shields et al. 2002). Totally there are 164 objects in our present sample.

Figure 1 shows the $v_{\text{FWHM}}(\text{H}\beta) - M_{\text{BH}}$ plot of the 164 AGNs. According to $L/L_{\text{Edd}} = \xi L_{5100}/L_{\text{Edd}}$ and $\xi = 5$, 19 of 84 quasars in Shields et al.'s sample are super-Eddington accretors. The

TABLE 1
LIST OF CANDIDATES OF SUPER-EDDINGTON AGNS

No.	Name	FWHM (H β)	$\log(M_{\text{BH}}/M_{\odot})$		Rev.	note
			$M_R - M_{\text{BH}}$	$\sigma - M_{\text{BH}}$		
1.	PG 0157+001	2140	9.19	...	8.18	DM01
2.	Q 0204+292	1040	8.69	...	7.13*	DM01
3.	PG 1001+054	1740	...	8.77	7.65	S02
4.	PG 1351+640	1170	...	...	7.66	K00
5.	PG 1440+356	1450	...	8.12	7.33	S02
6.	Q 2247+140	2220	8.94	...	8.07	DM01
7.	MS 2254-37	1545	...	8.17	7.04	S02

NOTES— S02: Shields et al. (2002); DM01: McLure & Dunlop (2001); K00: Kaspi et al. (2000).

* there is an error in estimation of 0204+292 in DM01.

** FWHM(H β) is in unit of km s⁻¹.

values of $\log L/L_{\text{Edd}}$ of the 19 quasars span from zero (VCV 0331-37) to the maximum 0.7 (PG 1216+069), however these objects are located far below the Eddington limit line as shown by Figure 1. If we take $\xi = 9$, there will be a quite large fraction of the objects with $L/L_{\text{Edd}} > 1$ and most of these objects are still located far below the Eddington limit lines. It follows from the Eddington limit that these objects have sub-Eddington rates though they have $L/L_{\text{Edd}} > 1$. We find that PG 1001+054, Mrk 478 and MS 2254-37 with $L/L_{\text{Edd}} < 1$ are located above the Eddington limit line. The three may be super-Eddington accretors. This inconsistency between the Eddington limit line and $\xi L_{5100}/L_{\text{Edd}}$ values in some objects reflects: 1) the intrinsic scatter of ξ , namely, the ratio $\xi L_{5100}/L_{\text{Edd}}$ is not fully reliable to estimate L/L_{Edd} as noted by Netzer (2003); 2) the estimation of the black hole mass via FWHM([O III]) has a larger uncertainty. As shown by Boroson (2003) in SDSS EDR sample of 107 radio-quiet quasars and Seyfert 1 galaxies, the correlation between FWHM([O III]) and M_{BH} is rather scatter, where M_{BH} is determined by Kaspi et al.'s empirical relation of reverberation mapping. The FWHM(H β) - M_{BH} relation also indicates the scatters of FWHM([O III]) - M_{BH} relation by this inconsistency. Therefore, the FWHM(H β) - M_{BH} relation may be more reliable to test whether an object has a super-Eddington accretion rate if the mass of the black hole is well determined. Mrk 478, PG 1001+054 and MS 2254-37 are narrow line Seyfert 1 galaxies [FWHM(H β) < 2000 km s⁻¹] (Grupe et al. 1999). Table 1 lists the estimation of the black hole masses of the three objects. We find that the differences of M_{BH} given by $\sigma - M_{\text{BH}}$ and the empirical relation of reverberation mapping are large, 1.1, 0.8 and 1.1 dex for PG 1001+054, Mrk 478 and MS 2254-37, respectively, showing the mass given by $\sigma - M_{\text{BH}}$ is larger than that given by the empirical relation of reverberation mapping. Even we use the masses by empirical relation of reverberation mapping, the three objects are still located on or close to the Eddington limit line. We thus suggest that they are probably super-Eddington accretors.

In the sample of McLure & Dunlop (2001), we take the data of Mrk 355 from the reverberation mapping measurements by Kaspi et al. (2000) and the data of $M_{\text{BH}} = 1.4^{+1.0}_{-0.6} \times 10^6 M_{\odot}$ and FWHM(H β) = 1110 $\pm$ 190 km s⁻¹ in NGC 4051 from Peterson et al. (2000). The values of L/L_{Edd} are given from L_{5100} , object with $L/L_{\text{Edd}} > 1$ has not been found in this sample. But the three radio-quiet quasars PG 0157+001, Q 0204+292 and Q 2247+140 have been found above the Eddington limit line. Table 1. also lists the three objects. We find that the masses of the black holes in PG 0157+001, Q 0204+292 and Q 2247+140 are quite different from $M_R - M_{\text{BH}}$ and reverberation relation methods. The mass of the black hole in Q 0204+292 is of $10^{8.9} M_{\odot}$ from $M_R - M_{\text{BH}}$ relation since the magnitude of its host galaxy is quite bright $M_R = -23.3$ (McLure & Dunlop 2001). On the other hand its FWHM(H β) = 1040 km s⁻¹ is very narrow, the

black hole mass is $10^{7.13} M_{\odot}$. The three objects locate above the Eddington limit lines even for their masses from empirical reverberation relation. We suggest that the three objects may be candidates of super-Eddington accretors.

No object with $L/L_{\text{Edd}} > 1$ has been found in Kaspi et al.'s sample, but PG 1351+640 locates above Eddington limit line. It may thus be a super-Eddington accretor. No super-Eddington has been found in the radio-loud quasar sample of Wang, Ho & Staubert (2002), who obtain the Eddington ratio from the converting the total emission line luminosity into ionizing luminosity. There is only one marginal object, PG 2201+315, which is located close to the Eddington limit whereas the others locate below the Eddington limit. We would like to stress that the above remarks are based on that the black hole masses are fully reliable.

For Kaspi et al.'s relation $\beta = 0.7$, we find a handful of objects that locate above the Eddington limit lines. These objects may be really super-Eddington accretors. For $\beta = 0.58$ empirical relation (Netzer 2003), we find that the FWHM(H β) - M_{BH} relation becomes flatter ($M_{\text{BH}} \propto v_{\text{FWHM}}^{5.0}$) and there are 23 of 164 ($\approx 14\%$) objects that locate above the Eddington limit lines (but they are not far away from the Eddington limit lines). If the black hole masses of these objects are less uncertain, they are likely super-Eddington accretors. We note that these objects locate the region between the Begelman and Eddington limit lines. One interesting possibility is that these objects are undergoing a hybrid process of photon trapping and photon bubble instability. It would be necessary to do the observation of reverberation mapping for these objects or measure the stellar dispersion velocity of their host galaxies in order to more reliably determine the black hole masses and the structure of the super-Eddington accretion disks. Additionally, the variabilities in the X-ray band will be very helpful to understand the physics in these objects (photon trapping or photon bubble instability).

All the objects listed in Table 1 have a narrower width of H β (< 2300 km s⁻¹). Based on the limit relation, one of prominent features of super-Eddington accretion candidates is the presence of the relatively narrow width of H β . If the largest black hole mass is of $M_{\text{BH}}^{\text{max}} = 2 \times 10^{10} M_{\odot}$ among ANGs (Netzer 2003), the maximum Eddington width is given by

$$V_{\text{FWHM}}^{\text{max}}(\text{H}\beta) = (3.0 - 3.8) \times 10^3 \left(\frac{M_{\text{BH}}^{\text{max}}}{2.0 \times 10^{10} M_{\odot}} \right)^{0.15} \text{ km s}^{-1}. \quad (7)$$

Therefore searching super-Eddington AGNs should be confined in those with H β width narrower than this Eddington width $V_{\text{FWHM}}^{\text{max}}(\text{H}\beta) = 3.0 - 3.8 \times 10^3 \text{ km s}^{-1}$. An interesting work is to apply the present limits to narrow line Seyfert 1 galaxies, which are regarded to be less massive black hole with higher accretion rates. The instability may make the disk inhomogeneous (Fabian et al. 2002). However the current data does not allow us to do so. It is suggested to do measurements of reverberation mapping for a sample of NLS1 for the determination of the black hole masses in order to probe the structures of the super-Eddington accretion disks.

The objects in the present samples used in this paper are low redshift. The above discussion can be easily extended to the relation between C IV/Mg II width and the mass of the black hole for high redshift quasars using the empirical relation of reverberation mapping (Vestergaard 2002, Netzer 2003).

Finally, we would like to point out the response of BLR size to the increases of disk's luminosity. According to the Kaspi et al.'s relation, the virializing timescale of the clouds in BLR is

of $\tau_{\text{vir}} = R_{\text{BLR}}/v_{\text{FWHM}} \approx 7.0 M_{\text{BH},8}^{-1/2} (L_{5100}/10^{44} \text{ erg s}^{-1})^{1.05}$ (yrs), where $M_{\text{BH},8} = M_{\text{BH}}/10^8 M_{\odot}$, which is much shorter than the super-Eddington accretion timescale ($\sim 10^6$ yrs). Thus the locations of $H\beta$ -emitting clouds will be rapidly re-arranged in BLR according to the luminosity. If the super-Eddington accretion lasts several million years, we should observe its effects on the width of broad emission lines.

4. CONCLUSIONS

Based on the super-Eddington accretion disk models of the photon trapping and photon bubble instability, we have derived a limit relation between the black hole mass and $H\beta$ width via the empirical relation of reverberation mapping in AGNs. We apply the present limits to several samples. We find that most of the objects locate below the Eddington limit line. In Shields et al.'s sample, there is a significant fraction of the objects have $\xi L_{5100}/L_{\text{Edd}} > 1$, but they are far below the Eddington limit line. This inconsistency implies that the present

limit relation is more reliable and convenient to test the super-Eddington accretions in AGNs. We find that there are a handful of objects locate above the Eddington limit lines, but below the Begelman limit line. We suggest they may be candidates of super-Eddington accretors. The maximum width of $H\beta$ is $3.0 - 3.8 \times 10^3 \text{ km s}^{-1}$ for the maximum black holes with super-Eddington accretion rates among AGNs, which confines the candidates in future search for super-Eddington accretions among AGNs. The $\text{FWHM}(H\beta) - M_{\text{BH}}$ relation is more reliable and convenient to test whether the objects have super-Eddington accretion rates, and can also be used to probe the structure of the super-Eddington accretion disks in AGNs.

The helpful comments from an anonymous referee are thanked. I am very grateful to Hagai Netzer for numerous useful suggestions and comments on the manuscript. The supports from Alexander von Humboldt Foundation, the Special Funds for Major State Basic Research Projects and NSFC are acknowledged.

REFERENCES

- Abramowicz, M.A., Czerny, B., Lasota, J.P., & Szuszkiewicz, E., 1988, *ApJ*, 322, 646
 Begelman, M. C., 1978, *MNRAS*, 184, 53
 Begelman, M. C., 2001, *ApJ*, 551, 897
 Begelman, M. C., 2002, *ApJ*, 568, L97
 Begelman, M. C. & Meier, D. L., 1982, *ApJ*, 253, 873
 Boroson, T., 2002, *ApJ*, 565, 78
 Boroson, T., 2003, *ApJ*, in press, astro-ph/0211372
 Boroson, T. & Green, R. F., 1992, *ApJS*, 80, 109
 Chen, X., et al., 1995, *ApJ*, 443, L61
 Chen, X. & Taam, R., 1993, *ApJ*, 412, 254
 Collin, S. & Huré, J.-M., 2001, *A&A*, 372, 50
 Collin, S., et al. 2002, *A&A*, 388, 771
 Dunlop, J.S., et al., 2001, astro-ph/0108397
 Fabian, A. C., et al., 2002, *MNRAS*, 331, L35
 Ferrarese, L. & Merritt, D., 2000, *ApJ*, 539, L9
 Fromerth, M.J. & Melia, F., 2000, *ApJ*, 533, 172
 Frank, J., King, A.R. & Raine, D.J., 2002, *Accretion Power in Astrophysics*, 2nd Edition (Cambridge University Press: Cambridge)
 Gammie, C. F., 1998, *MNRAS*, 297, 929
 Gebhardt, K., 2000, *ApJ*, 539, L13
 Grupe, D., et al., 1999, *A&A*, 350, 805
 Ho, L.C., 1999, in *Observational Evidence for Black Holes in the Universe*, ed. S.K. Chakrabarti (Dordrecht: Kluwer), 157
 Kaspi, S., et al., 2000, *ApJ*, 533, 631
 King, A., 2002, *MNRAS*, 335, L13
 King, A. & Puchnarewicz, E. M., 2002, *MNRAS*, 336, 445
 Kormendy, J. & Richstone, D., 1995, *ARAA*, 33, 581
 Kormendy, J. & Gebhardt, K., 2001, in *The 20th Texas Symposium on Relativistic Astrophysics*, ed. H Martel & J. C. Wheeler (Melville: AIP), 363
 Krolik, J. H., 2001, *ApJ*, 551, 72
 McLure, R.J. & Dunlop, J.S., 2001, *MNRAS*, 327, 199
 McLure, R.J. & Dunlop, J.S., 2002, *MNRAS*, 331, 795
 Mineshige, S., Kawakuchi, T., Takeuchi, M. & Hayashida, K., 2000, *PASJ*, 52, 499
 Nelson, C. H., 2000, *ApJ*, 544, L91
 Netzer, H., 2003, *ApJ*, 583, L3
 Netzer, H. & Laor, A., 1993, *ApJ*, 404, L51
 Ohsuga, K., Mineshige, S., Mori, M. & Umemura, M., 2002, *ApJ*, 574, 315
 Onken, C. A. & Peterson, B. M., 2002, *ApJ*, 572, 746
 Peterson, B. M., McHardy, I. M., et al., 2000, *ApJ*, 542, 161
 Peterson, B. M. & Wandel, A., 1999, *ApJ*, 521, L95
 Rees, J.J., 1984, *ARAA*, 22, 471
 Shakura, N. I., & Sunyaev, R. A. 1973, *A&A*, 24, 337
 Shields, G. A., Gebhardt, K., Salvander, S., Wills, B. J., Bingrong, X., Brotherton, M. S., Yuan, J., & Dietrich, M., 2002, *ApJ*, 583, 124
 Telfer, R. C., Zheng, W., Kriss, G. A. & Davidsen, A.F., 2002, *ApJ*, 565, 773
 Tremaine, S., et al., 2002, *ApJ*, 574, 740
 Vestergaard, M., 2002, *ApJ*, 571, 733
 Wandel, A., Peterson, B. M., & Malkan, M. A., 1999, *ApJ*, 526, 579
 Wang, J.-M., Ho, L. C. & Staubert, R., 2002, *A&A*, submitted
 Wang, J.-M. & Netzer, H., 2003, *A&A*, 398, 927
 Wang, J.-M., Szuszkiewicz, E., Lu, F.J. & Zhou, Y.Y., 1999, *ApJ*, 522, 839
 Wang, J.-M., & Zhou, Y.-Y., 1999, *ApJ*, 516, 420